

Emergent spin excitations in a Bethe lattice at percolation

Hitesh J. Changlani,¹ Shivam Ghosh,¹ Sumiran Pujari,² and Christopher L. Henley¹

¹Laboratory of Atomic And Solid State Physics, Cornell University, Ithaca, NY 14853, USA

²Laboratoire de Physique Théorique, Université Paul Sabatier, 31062 Toulouse, France

(Dated: October 16 2012)

We study the spin 1/2 quantum Heisenberg antiferromagnet on a Bethe lattice diluted to the percolation threshold. Dilution creates areas of even/odd sublattice imbalance resulting in “dangling spins” (L. Wang and A. W. Sandvik, Phys. Rev. Lett. 97, 117204 (2006); Phys. Rev. B 81, 054417 (2010)). These collectively act as “emergent” spin 1/2 degrees of freedom and are responsible for the creation of a set of low lying “quasi degenerate states”. Using Density Matrix Renormalization Group (DMRG) calculations, we detect the presence and location of these emergent spins. We find an effective Hamiltonian of these emergent spins, with Heisenberg interactions that decay exponentially with the distance between them.

Introduction—Quantum spins on percolation clusters [1] provide an ideal testbed for studying the interplay between geometrical disorder and quantum fluctuations. The Hamiltonian for these problems is

$$\mathcal{H} = \sum_{\langle ij \rangle} J \mathbf{S}_i \cdot \mathbf{S}_j \quad (1)$$

where $\mathbf{S}_i$ are Pauli spin 1/2 operators and the sum runs over nearest-neighbor occupied sites, and $J > 0$. Theoretical [2, 3] and experimental [4] studies of quantum spins on diluted square lattices have focussed on the question of whether long range order survives up to the classical percolation threshold p_c . A numerical study [5] seems to have settled this question and found long range order to be robust to quantum fluctuations, surviving all the way up to p_c .

The excitations are less straightforward. For uniform lattices with number of sites N , the lowest energy scale consistent with Néel order breaking a continuous symmetry is $\sim JN^{-1}$, corresponding to a “tower” of states: mixtures of symmetry-broken states that become degenerate in the thermodynamic limit [6–8]. However a Quantum Monte Carlo study by Wang and Sandvik [9] discovered a somewhat “anomalous” finite size scaling of the spin gap Δ_{low} : $\Delta_{low} \approx N^{-2}$ (for clusters with a singlet ground state) or $\Delta_{low} \approx N^{-1.5}$ (for generic clusters, most with a non-singlet ground state). A strong correspondence was shown [9] between these low lying states and places on the cluster where there is a local imbalance between the number of even and odd sites. It was conjectured that, in each such place, a spin degree emerges which is effectively decoupled from the antiferromagnetic order and hence was called a “dangling spin.”

The goal of this Letter is to characterize the dangling-spin degrees of freedom numerically, relating their nature to the local geometry of the cluster, and to explain the observed low energy spectrum in terms of mediated interactions between dangling spins. Our Hamiltonian is (1) on clusters obtained by randomly diluting the Bethe Lattice of coordination 3 at its percolation threshold, $p_c = 1/2$ (see examples of small clusters in Fig. 1). The lack of loops in the Bethe lattice is conducive for using the Density Matrix Renormalization Group (DMRG) algorithm [10], as adapted to generic tree graphs, to obtain ground and (some) excited states [11].

In the rest of this Letter, we first show that a typical percolation cluster’s spectrum has a clearly separated low energy




Figure 1: (Color online) Three different percolation clusters (all of the same size $N = 18$) are shown with their corresponding low energy spectra. The ‘red’ and ‘green’ circles indicate even and odd sites. The broken dashed lines show dimer coverings which serve as a heuristic to locate the “dangling spins”. Dangling spins in each cluster have been circled with thick black lines. Low energy spectra for each of the clusters show low lying quasi degenerate (QD) states separated from the continuum by an energy scale Δ . σ_{QD} (not shown) is a measure of the spread of energies in the QD space.

component, with a multiplicity consistent with the expected number of weakly coupled spin-1/2 (sometimes spin-1) dangling spins. We next show that each dangling spin is somewhat delocalized over a few sites: on the one hand, we model it as an unpaired spin in a dimerized background to predict the dangling spin’s nature from the local geometry; on the other hand, by processing spin expectations we obtain the explicit “localized state wavefunction” for each dangling spin. Finally, for each cluster we construct the effective Hamiltonian of the emergent dangling spins, consisting of *pairwise*, unfrustrated exchange interactions decaying exponentially with separation, mediated by the background of almost dimerized spins on the balanced parts of the cluster; this accurately reproduces the details of that cluster’s low-energy spectrum.

Exact correspondence between dangling spins and low energy spectrum— We carried out DMRG calculations for several hundred balanced clusters (i.e. having equal number of

even and odd sublattice sites) [12] for sizes up to $N = 100$, targeting multiple excited states in the low energy spectrum. Since the number of low energy states was found to increase rapidly with an increase in the number of dangling spins, we restricted our analysis to the case of four dangling spins [13]. For analyses where only the ground state properties were of interest or where only the lowest energy gap Δ_{low} was required, larger clusters ($N = 200$), with an *arbitrary* number of dangling spins, were also studied.

In a typical percolation cluster, we observed a distinct set of low-lying energy levels we shall call "quasi-degenerate" (QD) [14] since (we claim) they would be exactly degenerate in the limit that the dangling spins are fully decoupled from the rest of the sites. The QD states are separated from the continuum of higher energy Quantum Rotor and spin wave states by a finite size gap we call Δ (specifically defined as the difference between the mean of QD levels and the lowest non-QD level). The set of QD states are identified by looking at the difference in energies of consecutive states up to the Quantum rotor excitation and finding the pair of states with the largest gap. The lower energy state in this pair and all states below that make up the QD spectrum [15]. The energy scale characterizing the spread of the QD states, σ_{QD} , is defined to be the standard deviation of the QD energies from their mean value. The ratio $r = \sigma_{QD}/\Delta$ was found to be small (for example $r \sim 0.17 \pm 0.1$ for $N = 50$), justifying our notion of a separation of scales.

Figure 1 also shows a striking correspondence between the number of low lying QD states, $\mathcal{N}_{QD}$, and the number of dangling spins n_d on the percolation cluster. We find that $\mathcal{N}_{QD} = 2^{n_{1/2}} 3^{n_1}$, where $n_{1/2}$ and n_1 are integers and $n_{1/2} + 2n_1 = n_d$. Our interpretation of this multiplicity is that $2n_1$ of the dangling spins pair up so as to form a spin-1, while the others remain as spin-half degrees of freedom. There is thus a one-to-one correspondence between the low-energy (QD) eigenstates and the Hilbert space of the posited emergent spins. We used an algorithm (to be described in the next section) that relies only on the cluster geometry to objectively predict the numbers $n_{1/2}$ and n_1 for each cluster, and verified that their predicted relationship with $\mathcal{N}_{QD}$ was satisfied in every cluster.

We also directly measured the lowest singlet-triplet gaps Δ_{low} for an ensemble of balanced clusters. Its typical value scales as $N^{-1.9 \pm 0.1}$, which appears remarkably similar to the scaling previously seen on square lattice percolation clusters [9].

Locating Dangling degrees of freedom in real space— Having established the presence of emergent spin-half and spin-one degrees of freedom, we now develop two complementary ways of looking at them.

The first is within the framework of a quantum monomer-dimer model. We imagine that the wavefunction is a product of valence bonds in which the N spins are paired (dimerized) into singlets to the maximum extent possible (optimal configuration). Even when even and odd sites are balanced *globally*, there remain some uncovered sites, i.e. monomers, due to *local* imbalances. These are spin-1/2 degrees of freedom and (within this picture) represent the dangling spins. There are

multiple ways to optimally place the monomers; the actual wavefunction is imagined to be a superposition of these ways.

Our geometric algorithm, based on the valence bond framework, finds one element from the set of optimal dimerizations of the cluster and then attempts to find other elements of the set by locally replacing monomers with adjacent dimers. In spirit, this is a "greedy" algorithm which tries to place dimers wherever possible (to obtain an optimal dimerization pattern), working from the outer sites inwards on the cluster.




Figure 2: (Color online): Typical geometrical motifs in Cayley tree percolation clusters, as related to monomer/dimer construction. We disconnect the cluster into "spans", shown by (blue) dashed circles, at the "prong" bonds, as indicated by the (red) cut lines. The thickness of the (grey) bonds is directly proportional to $\langle \mathbf{S}_i \cdot \mathbf{S}_j \rangle$.




Figure 3: (Color online): Spatial profiles associated with "dangling spins", shown on the subclusters marked (a)-(d) of the same cluster shown in Fig. 2. The area of the black circles is proportional to ω_{ii} given by Eq. (4). (a) shows a "fork" (b) shows a site surrounded by two "hairs" (c) shows a subcluster where two monomers on the same sublattice are present forming an effective spin-1. The (blue) arrows indicate the monomer is free to hop around (delocalize) within the subcluster. (d) shows a region where the spins are "inert" (largely dimerized) and do not participate in the low energy physics.

Given any cluster, there are two operations which cut it down to a smaller cluster or clusters, such that all optimal dimerizations on the smaller cluster(s) are in 1-to-1 correspondence with *some* of dimerizations on the larger one. The first operation is that, wherever two sites have coordination 1 or 2, we can remove both (given the dimerization on the smaller cluster, just insert another dimer to get the dimerization on the larger one). The second operation is that, wherever we find a pair of adjacent sites with respective coordinations 3 and 1 (a "prong"), we can always place a dimer on that pair,

which fragments the rest into two subclusters (Fig. 2); a very common special case is the fork, at which we can arbitrarily choose either side to be the “prong”. These two operations can be used recursively till only isolated sites remain, each corresponding to one monomer in the original cluster. Furthermore, any other optimal dimerization is accessible from the special one we constructed, by repeatedly exchanging a monomer with an adjacent dimer.

A monomer can thus “hop” to sites on the lattice via such local monomer-dimer rearrangements as shown in Fig. 3(c). Our rule-of-thumb is that two monomers (of the same sublattice) form a spin-1 if and only if they can approach to the minimal separation of two steps [16]. We check this possibility by enumerating all dimerizations accessible from the one generated by our algorithm.

Fig. 2 also shows some typical geometrical motifs seen in an ensemble of percolation clusters. In Fig. 2(a), a spin-half dangling spin is localized at the tip of the fork. The arms of the fork have a strong dimerization and prevent the localized monomer from hopping to other sites in the cluster. Fig. 2(b) shows another type of spin-half degree of freedom, which is trapped in the interval between (or on) two “prongs”. Fig. 2(c) is an example of an emergent spin-one excitation. For comparison, we also show in Fig. 2(d) a region of strong dimerization with no dangling spins.

Our second way to capture the spatial nature of a dangling spin degree of freedom starts from the idea that it could be adiabatically connected to an uncoupled spin, analogous to the Landau quasiparticle that is adiabatically connected to a free electron. Thus, our program is to label each emergent spin-1/2 degree of freedom by a “quasi spin” operator $\mathbf{T}_\alpha$, where the index α labels each region on the cluster with a local spin imbalance. The $\mathbf{T}_\alpha$ ’s are idealized as having a spin-half algebra. The actual “quasi spin” excitation is a composite object involving multi-spin excitations, localized on a few sites.

Our assumption is that the quasi-spin quantum numbers are sufficient to label all the QD states; furthermore, we expect the action of any spin operator $\mathbf{S}_i$, when restricted to the QD states, practically reduces to a linear combination of $\mathbf{T}_\alpha$ ’s acting on the quasi spins. Specifically, let $\hat{P}_{\text{QD}}$ be the projection operator onto the QD subspace. Let $\mathbf{S}_{i,\text{QD}} \equiv \hat{P}_{\text{QD}} \mathbf{S}_i \hat{P}_{\text{QD}}$. Then,

$$\mathbf{S}_{i,\text{QD}} \cong \sum_{\alpha} u_i^{(\alpha)} \mathbf{T}_{\alpha} \quad (2)$$

where each mode $u_i^{(\alpha)}$ has most of its weight on sites within the region α and is expected to decay rapidly outside [17, 18].

Two operators $\hat{P}, \hat{Q}$ are said to be *orthogonal* when their Frobenius inner product $(\hat{P}, \hat{Q})_F \equiv \text{Tr}(\hat{P}^\dagger \hat{Q})$ is exactly zero. In this sense, the $\mathbf{T}_\alpha$ operators are orthogonal to each other. Since each $\mathbf{T}_\alpha$ is a quasi-spin 1/2 operator its inner product with itself is 1/2 (for each spin component).

In light of Eq. (2), we can also construct a good approximation $\tilde{\mathbf{T}}_\alpha$ to each operator $\mathbf{T}_\alpha$, by choosing any representative site i in the dangling region α and normalizing the restriction

of its spin operator to the QD states:

$$\tilde{\mathbf{T}}_\alpha \equiv \frac{\mathbf{S}_{i,\text{QD}}}{\sqrt{2} \|\mathbf{S}_{i,\text{QD}}\|} \quad (3)$$

where $\|\hat{O}\| \equiv (\hat{O}, \hat{O})_F^{1/2}$ is the norm of any operator $\hat{O}$. Note that the $\tilde{\mathbf{T}}_\alpha$ ’s are *not* orthogonal to each other. A procedure to construct the $\mathbf{T}_\alpha$ ’s from the $\tilde{\mathbf{T}}_\alpha$ ’s will be discussed in the next section.

Given the proposed relationship of the bare spins to the quasi spins, we discuss two related but independent measurements to recover the mode vectors $u_i^{(\alpha)}$ from numerically evaluated expectations. For all the measurements proposed here, it was sufficient to record (within DMRG) all matrix elements of the form $\langle l' | \mathbf{S}_i^\pm | l \rangle$ and energies E_l for eigenstates l, l' in the QD subspace.

First, we consider the operator overlap ω_{ij} between two spins i and j on the lattice, defined to be,

$$\omega_{ij} \equiv (S_{i,\text{QD}}^+, S_{j,\text{QD}}^+)_F \quad (4)$$

We substitute our ansatz (2) into (4) and use the operator orthogonality of the $\mathbf{T}_\alpha$ ’s, to get $\omega_{ij} = \sum_{\alpha} u_i^{(\alpha)} u_j^{(\alpha)}$. If we consider a site i to be well within a dangling region α (i.e. $u_i^{(\alpha)}$ is relatively large) then the amplitude on the remaining sites j (but far away from other dangling regions) is approximately $u_j^{(\alpha)}$. Thus, the relative amplitudes of the mode vector can be recovered by this method.

Our second measurement involves computation of the inter-site spin susceptibility matrix,

$$\chi_{ij} = \int_0^\infty \langle \hat{S}_i^z(\tau) \hat{S}_j^z(0) \rangle_{GS} d\tau = \sum_n \frac{\langle 0 | \hat{S}_i^z | n \rangle \langle n | \hat{S}_j^z | 0 \rangle}{E_n - E_0} \quad (5)$$

where τ is imaginary time, $|0\rangle$ denotes the ground state and E_n is the energy of an excited state $|n\rangle$ [19].

Though the sum runs over all excited states, it can be well approximated by taking only the states in the QD subspace. Then χ_{ij} can also be expressed in terms of the mode profiles $u_i^{(\alpha)}$,

$$\chi_{ij} = \sum_{\alpha\beta} u_i^{(\alpha)} u_j^{(\beta)} X_{\alpha\beta}; \quad X_{\alpha\beta} \equiv \sum_{n \in \text{QD}} \frac{\langle 0 | \mathbf{T}_\alpha^z | n \rangle \langle n | \mathbf{T}_\beta^z | 0 \rangle}{E_n - E_0} \quad (6)$$

Consider site $i(j)$ in dangling region $\alpha(\beta)$. From equation (6) it follows that $\chi_{ij} \approx u_i^{(\alpha)} u_j^{(\beta)} X_{\alpha\beta}$, where the last factor is *independent* of sites i, j (so long as we stay within those regions). Within this approximation, the susceptibility matrix breaks up into blocks of rank 1 from which we can immediately pull out the $u_i^{(\alpha)}$ and $u_j^{(\beta)}$ modes.

Effective Hamiltonian in the Quasi degenerate subspace—According to our ansatz (2), there is a one-to-one correspondence between the QD Hilbert space and the Hilbert space of a set of abstract “quasi spin” operators $\mathbf{T}_\alpha$. (For simplicity, assume they all have spin 1/2.) The latter are labeled using an Ising basis $|\phi_t\rangle$, where t stands for the quantum numbers $\{t_1^z, t_2^z, \dots, t_{n_d}^z\}$, with $t_\alpha = \pm 1/2$. We want to find the

unitary matrix $\mathbf{M}$ of coefficients expressing the QD states $|l\rangle$ (in eigenenergy basis) in terms of the quasi-spin basis, $|l\rangle = \sum_t M_{lt} |\phi_t\rangle$.

Using $\tilde{\mathbf{T}}_\alpha$ from (3), we define $\hat{Q}_\alpha^{\pm 1/2} \equiv (\frac{1}{2} \pm \tilde{\mathbf{T}}_\alpha^z)$, which is *almost* a projection operator, and let

$$|\tilde{\phi}_t\rangle \propto \prod_{\alpha=1}^{n_d} \hat{Q}_\alpha^{t_\alpha} |n\rangle \quad (7)$$

where $|n\rangle$ could be any QD state (that is not annihilated by the operator prefactors) and $|\tilde{\phi}_t\rangle$ is normalized. Finally, define a matrix $\underline{\Omega}$ by $\Omega_{tt'} \equiv \langle \tilde{\phi}_t | \tilde{\phi}_{t'} \rangle$ – which is *almost* the identity matrix – and construct the orthonormal quasi spin basis of the QD states as

$$|\phi_t\rangle \equiv \sum_{t'} (\underline{\Omega}^{-1/2})_{tt'} |\tilde{\phi}_{t'}\rangle. \quad (8)$$

The quasi spin operators $\mathbf{T}_\alpha$ are then defined so as to have the appropriate matrix elements in this basis.

Now consider the effective low energy Hamiltonian written in terms of the many body eigenstates $|l\rangle$,

$$\mathcal{H}_{\text{eff}} \equiv \sum_{l \in \text{QD}} E_l |l\rangle \langle l| = \sum_{tt'} h_{tt'} |\phi_t\rangle \langle \phi_{t'}|, \quad (9)$$

where E_l is the eigenenergy of QD state $|l\rangle$, and the matrix elements $h_{tt'}$ can be calculated since we know the transformation between the bases $\{|l\rangle\}$ and $\{|\phi_t\rangle\}$. Every term $|\phi_t\rangle \langle \phi_{t'}|$ can be uniquely expressed as a polynomial in the spin operators $\{T_\alpha^z\}$ and $\{T_\alpha^\pm\}$.

The effective Hamiltonian (9) then takes the new form of,

$$\mathcal{H}_{\text{eff}} \equiv \sum_{\mu, \nu} J_{\mu\nu} \mathbf{T}_\mu \cdot \mathbf{T}_\nu + \text{multi spin terms}; \quad (10)$$

(The two-spin terms must have this Heisenberg form due to the exact rotational symmetry retained by the QD states.)

Although the magnitude of $J_{\mu\nu}$ depends on the detailed geometry of the cluster along the path connecting dangling regions μ and ν , roughly speaking it decays with distance (using as metric the number of steps within the Bethe lattice, the so-called “chemical” distance). This is quantified by the scatter plots in Fig. 4, for an ensemble limited to clusters of equal size $N = 100$ [20] each having two dangling spin-1/2 spins ($n_d = n_{1/2} = 2$). Since each dangling region is in general spread out over multiple sites, we must define an “effective distance” between two of them:

$$\bar{d}_{\mu\nu} \equiv \sum_{ij} |u_i^{(\mu)}|^2 |u_j^{(\nu)}|^2 d_{ij} \quad (11)$$

where d_{ij} is the distance between sites i and j belonging to dangling regions μ and ν , and the amplitudes $u_i^{(\alpha)}$ for mode α are normalized. Figure 4 shows that indeed $J_{\mu\nu} \approx J_0 e^{-\bar{d}_{\mu\nu}/\xi}$, where $(J_0, \xi) \approx (+0.15, 4.85)$ for an even/odd pair of dangling spins, which are *always* antiferromagnetically coupled, or $(-0.22, 4.55)$ for a pair on the same sublattice. In




Figure 4: (Color online) Effective couplings J_{12} between two dangling spins as a function of their effective separation obtained from clusters having exactly two dangling spins. (a) dangling spins on opposite sublattices (antiferromagnetic coupling) (b) on same sublattice (ferromagnetic coupling).

the *ferromagnetic* case, choosing to fit only to clusters which do not form a spin-1 gives parameter values closer to the *antiferromagnetic* case.

We considered another ensemble (not plotted) of clusters with $N = 50$ having four dangling spins ($n_d = 4$) and obtained the effective Hamiltonian using the same prescription. We found that the non-pair terms typically account for a weight of at most 5% (using the Frobenius norm) in the Hamiltonian (10), confirming that the effective Hamiltonian is well approximated by pairwise Heisenberg exchange (at least in the limit of dilute monomer concentration).

Conclusion— We addressed the excitations of the spin 1/2 Heisenberg antiferromagnet on Bethe lattice percolation clusters. The emergent low energy degrees of freedom were found to be composite “dangling spin” excitations arising wherever there is a local imbalance between the even and odd sublattices [9]. (The same kind of imbalance was also recently shown to determine the low energy spectra of *regular* Cayley trees [11].) We classified dangling spins based on the local cluster geometry and described an algorithm to predict their locations on clusters of arbitrary size.

Each of these emergent degrees of freedom is associated with a profile $(u_i^{(\alpha)})$ in the text that plays the role of a “spinon wavepacket wavefunction” [21]; the degree of localization of the dangling spins could, in future work, be quantified using metrics (e.g. localization lengths) borrowed from Anderson localization. We did not, however, address the fundamental

reason why a dangling spin degree of freedom decouples from the cluster, an effectively zero-energy excitation; that will be addressed in a subsequent publication [17].

The dangling spins interact via an effective low energy Hamiltonian, consisting of unfrustrated Heisenberg couplings that decay (roughly) exponentially with separation. If we adopt the fitted dependences in Figure 4 as our *definition* of our Hamiltonian, it should be possible to understand the low-energy states for arbitrary clusters that are much too large for the DMRG (involving thousands of sites), but will be tractable using the strong disorder renormalization group method [22], and thus finally explain the scaling of the spin gap Δ_{low} with cluster size that was found in Refs. 9. It would be interesting to see if the finite size scaling is *exactly* N^{-2} and if so,

whether it is universal.

Finally, how does long range magnetic order survive in the scenario of dangling spins mediating interactions across strongly dimerized inert regions? We observed that spin correlations had an exponential decay across dimers but revived in regions on the cluster where a dangling spin was located. This suggests that in the thermodynamic limit, locally unbalanced regions are crucial for the propagation of magnetic order on percolation clusters.

Acknowledgement—HJC would like to thank Garnet Chan for discussions on the DMRG technique. We also thank Anders Sandvik for discussions and Andreas Läuchli for pointing out references on related work. We acknowledge support from National Science Foundation grant NSF DMR-1005466.

-
- [1] D. Stauffer and A. Aharony, *Introduction to Percolation Theory* (Taylor and Francis, New York, 1992).
 - [2] Y.-C. Chen and A. H. Castro Neto, Phys. Rev. B 61, R3772 (2000).
 - [3] N. Bray-Ali, J. E. Moore, T. Senthil and A. Vishwanath, Phys. Rev. B. 73, 064417 (2006).
 - [4] O. P. Vajk, P.K. Mang, M. Greven, P.M. Gehring and J.W. Lynn, Science 295, 1691 (2002).
 - [5] A. W. Sandvik, Phys. Rev.B 66, 024418 (2002).
 - [6] P.W. Anderson, Phys. Rev. 86, 694 (1952), P.W. Anderson, *Basic Notions of Condensed Matter Physics* (Benjamin, New York, 1984), see section 2D, Page 44-46.
 - [7] H. Neuberger and T. Ziman, Phys. Rev. B 39, 2608 (1989).
 - [8] M. Gross, E. Sanchez-Velasco and E.Siggia, Phys. Rev. B 39, 2484 (1989).
 - [9] L. Wang, A. W. Sandvik, Phys. Rev. Lett. 97, 117204 (2006), L. Wang and A. W. Sandvik, Phys. Rev. B 81, 054417 (2010).
 - [10] S.R. White, Phys. Rev. Lett. 69, 2863 (1992).
 - [11] H. J. Changlani, S. Ghosh, C. L. Henley and A.M. Läuchli, arXiv:1208.1773.
 - [12] Balanced clusters necessarily have a singlet ground state.
 - [13] When only a count of the number of low energy states was desired, we looked at the approximate energies obtained at the end of the initialization step of the DMRG (as described in [11]). This allowed us to probe clusters with six dangling spins.
 - [14] C. Lhuillier arXiv:cond-mat/0502464v1 (unpublished).
 - [15] Interactions between dangling spins separated by different length scales creates an entire hierarchy of energy levels. Some of these might even be higher than the first Quantum Rotor excitation. However, there will always be a set of low lying states arising from splittings between the most distant spins on the cluster. We label these states as 'QD' spectrum for computing r .
 - [16] This prescription for predicting spin ones was found to be invalid for 3% clusters with $n_d = 4$. For these clusters, two monomers at lattice spacings greater than two, were found to behave as a spin one in the low energy spectrum. A more stringent criterion for forming a spin one is to have the relative spacing between the spin one monomers to be much smaller than the typical separation of the spin one pair from other spins on the cluster.
 - [17] S. Ghosh, H. J. Changlani and C. L. Henley (unpublished).
 - [18] The spatially localized mode u_i can also be obtained within Schwinger boson mean field theory and will be the subject of Ref. [17].
 - [19] The χ_{ij} matrix is a generalization of the χ_{ii} measurement proposed in Ref. 9.
 - [20] We observed a weak dependence of the coupling (for a given "effective distance") on the size of the cluster. A more detailed analysis will be carried out elsewhere.
 - [21] Y. Tang and A. W. Sandvik, Phys. Rev. Lett. 107,157201 (2011).
 - [22] R. N. Bhatt and P. A. Lee, Phys. Rev. Lett. 48, 344 (1982).